

# The Selective Na<sub>v</sub>1.7 Inhibitor, PF-05089771, Interacts Equivalently with Fast and Slow Inactivated Na<sub>v</sub>1.7 Channels

Jonathan W. Theile, Matthew D. Fuller, and Mark L. Chapman

Neusentis US, Pfizer Global R&D, (currently Icagen, Inc.), Durham, North Carolina

Received June 2, 2016; accepted August 31, 2016

## ABSTRACT

Voltage-gated sodium (Na<sub>v</sub>) channel inhibitors are used clinically as analgesics and local anesthetics. However, the absence of Na<sub>v</sub> channel isoform selectivity of current treatment options can result in adverse cardiac and central nervous system side effects, limiting their therapeutic utility. Human hereditary gain- or loss-of-pain disorders have demonstrated an essential role of Na<sub>v</sub>1.7 sodium channels in the sensation of pain, thus making this channel an attractive target for new pain therapies. We previously identified a novel, state-dependent human Na<sub>v</sub>1.7 selective inhibitor (PF-05089771, IC<sub>50</sub> = 11 nM) that interacts with the voltage-sensor domain (VSD) of domain IV. We further characterized the state-dependent interaction of PF-05089771 by systematically varying the voltage, frequency, and duration of conditioning prepulses to provide access to closed, open, and

fast- or slow-inactivated states. The current study demonstrates that PF-05089771 exhibits a slow onset of block that is depolarization and concentration dependent, with a similarly slow recovery from block. Furthermore, the onset of block by PF-05089771 develops with similar rates using protocols that bias channels into predominantly fast- or slow-inactivated states, suggesting that channel inhibition is less dependent on the availability of a particular inactivated state than the relative time that the channel is depolarized. Taken together, the inhibitory profile of PF-05089771 suggests that a conformational change in the domain IV VSD after depolarization is necessary and sufficient to reveal a high-affinity binding site with which PF-05089771 interacts, stabilizing the channel in a nonconducting conformation from which recovery is slow.

## Introduction

Voltage-gated sodium (Na<sub>v</sub>) channels play a key role in regulating action-potential generation and propagation in excitable cells (Hodgkin and Huxley, 1952). The Na<sub>v</sub> channel is a heteromeric protein complex consisting of a pore-forming  $\alpha$ -subunit and auxiliary  $\beta$ -subunits (Catterall, 2000). The  $\alpha$ -subunit consists of four homologous domains (DI–DIV), each containing six membrane spanning segments (S1–S6). The S1–S4 segments together comprise the voltage-sensing domain (VSD), in which translocation of the positively charged S4 segment after depolarization plays a critical role in the initiation of voltage-dependent gating (Noda et al., 1984; Stühmer et al., 1989; Catterall, 2010). The S5–S6 segments of each domain converge to form the ion conducting pore module, with the residues along the S6 segment constituting the inner pore (Ragsdale et al., 1994).

Inhibitors selective for peripherally expressed Na<sub>v</sub>1.7, Na<sub>v</sub>1.8, and Na<sub>v</sub>1.9 isoforms have been identified as potential analgesics for the treatment of pain (Dib-Hajj et al., 2009; Theile and Cummins, 2011; Payne et al., 2015), particularly Na<sub>v</sub>1.7 (Cox et al., 2006; Dib-Hajj et al., 2008;

Alexandrou et al., 2016). Current pharmacological treatments for pain that demonstrate activity against Na<sub>v</sub> channels include local anesthetics, tricyclic antidepressants, and anticonvulsants; however, many of these treatment options do not provide adequate pain relief and exhibit adverse central nervous system and cardiac toxicities, presumably due to a lack of pharmacological selectivity across the Na<sub>v</sub> family (Cummins and Rush, 2007). These classic Na<sub>v</sub> channel blockers bind to conserved residues along the pore-forming S6 segment of the Na<sub>v</sub> channel, resulting in little to no selectivity among Na<sub>v</sub> channel subtypes (Ragsdale et al., 1994, 1996; Yarov-Yarovsky et al., 2001, 2002). The typically narrow therapeutic index of classic Na<sub>v</sub> blockers is attributable to a degree of functional selectivity achieved through state- and/or frequency-dependent inhibition (Fozzard et al., 2005). Thus, there exists a need for the development of small molecule inhibitors with novel binding properties and Na<sub>v</sub> channel subtype selectivity to enhance clinical utility while minimizing undesired side effects. Because gating modifier toxins have been shown to affect normal voltage-dependent gating of Na<sub>v</sub> channels through an interaction with less conserved extracellular residues of the VSD and the nearby pore module (Catterall et al., 2007; Wang et al., 2011; Xiao et al., 2014), the development of small molecules that alter VSD function may display improved specificity and isoform selectivity over current treatment options.

We recently reported a class of low nanomolar-potency, state-dependent Na<sub>v</sub> inhibitors that bind to a novel site on the

This work was previously presented at the following workshop: Theile JW, Fuller MD, and Chapman ML (2015) NAV1.7 Inhibitor, PF-05089771, Inhibits Fast- and Slow-Inactivated Channels with Similar Affinities (2902-Pos/B332). *Biophysical Society Meeting*; February 7–11, 2015; Baltimore, MD.

This work was funded by Pfizer Ltd.  
dx.doi.org/10.1124/mol.116.105437.

**ABBREVIATIONS:** ANOVA, analysis of variance; DI–DIV, domains I–IV; DMSO, dimethyl sulfoxide; Na<sub>v</sub>, voltage-gated sodium channels; PM, pore module; S1–S6, transmembrane segments 1–6; VSD, voltage-sensing domain.

VSD of DIV, conferring a high degree of selectivity among Na<sub>v</sub> isoforms (McCormack et al., 2013). Within this class of aryl sulfonamide small molecules, PF-05089771 is a novel, selective, and potent human Na<sub>v</sub>1.7 inhibitor, exhibiting an IC<sub>50</sub> of ~11 nM for inactivated channels (Alexandrou et al., 2016). In addition to the potency and selectivity, PF-05089771 was reported to be distinguished by its strong preference for inactivated channels and the slow rate at which inhibition develops. In that initial report by Alexandrou and colleagues, it was observed that long applications of compound were required to achieve steady-state inhibition of Na<sub>v</sub>1.7 channels at concentrations near the IC<sub>50</sub>. One possible interpretation of this observation is that PF-05089771 exhibits a preference for a slow-inactivated state of the channel that accumulates with repeated activation. However the previous series of experiments did not directly address the possibility of slow association with multiple inactivated conformations. In this report, we sought to differentiate between these possibilities by altering the relative availability of the various kinetic states of the channel. We investigated this question using voltage-clamp protocols that biased channels into predominantly fast- or slow-inactivated state populations. We observed that inhibition occurs after depolarization of the channel, with the time course for inhibition largely unaffected by the relative availability of either open or kinetically distinct inactivated states. These results indicate that depolarization of the channel into a nonresting conformation is the primary determinant for PF-05089771-mediated inhibition, and any subsequent conformational changes associated with the progressive development of inactivation did not substantially alter the interaction of PF-05089771 with the channel.

## Materials and Methods

**Reagents.** PF-05089771 [2,2-diphenyl-*N*-(4-(*N*-thiazol-2-ylsulfamoyl)-phenyl)acetamide] was synthesized by the medicinal chemistry group at Neusentis, Durham, NC (Alexandrou et al., 2016).

**Cell Culture.** Human Na<sub>v</sub>1.7 was stably expressed in human embryonic kidney 293 cells. Methods of stable cell line generation were as described in McCormack et al. (2013).

**Electrophysiology.** Coverslips with human embryonic kidney 293 cells expressing human Na<sub>v</sub>1.7 were placed in a recording chamber on the stage of an inverted microscope and perfused with an extracellular solution containing (in mM), 132 NaCl, 1.8 CaCl<sub>2</sub>, 5.4 KCl, 0.8 MgCl<sub>2</sub>, 5 glucose, and 10 HEPES, pH 7.4, with NaOH. Recording patch pipettes were filled with an intracellular solution containing (in mM), 110 CsF, 35 CsCl, 5 NaCl, 10 EGTA, 10 HEPES, pH 7.3 with CsOH, and had a resistance of 1 to 3 MΩ. All reagents used for buffers were purchased from Sigma-Aldrich (St. Louis, MO). All recordings were made at room temperature (22–24°C) using Axopatch 200B or Multiclamp 700B amplifiers and PCLAMP software (Molecular Devices, Sunnyvale, CA). Sodium currents were measured using the whole-cell configuration of the patch-clamp technique. All compounds were dissolved in dimethyl sulfoxide (DMSO) to make 10 mM stock solutions, which were then diluted into extracellular solution to attain the final concentrations desired. The final concentration of DMSO (<0.1%) was found to have no significant effect on sodium currents. Assessment of recovery from inactivation and development of inhibition in the presence of PF-05089771 are described in the text and figure legends.

**Data Analysis.** Data were analyzed using the software programs of Clampfit 10.3 (Molecular Devices) and GraphPad Prism 6.0 (GraphPad Software, Inc., San Diego, CA). Statistical analysis was calculated by Student's *t* test or analysis of variance (ANOVA) where

mentioned, and  $P < 0.05$  indicates a significant difference. All data are presented as mean  $\pm$  S.E.M., and error bars in figures represent S.E.M.


Peak inward currents obtained from activation protocols were converted to conductance values using the equation,  $G = I/(V_m - E_{Na})$ , for which  $G$  is the conductance,  $I$  is the peak inward current,  $V_m$  is the membrane potential step used to elicit the response, and  $E_{Na}$  is the reversal potential for sodium (determined for each cell using the  $x$ -axis intercept of a linear fit of the peak inward current responses). Conductance data were normalized by the maximum conductance value and fit with a Boltzmann equation of the form  $G = G_{min} + (G_{max} - G_{min})/(1 + \exp[(V_{1/2} - V_m)/k])$ , where  $V_{1/2}$  is the midpoint of activation and  $k$  is a slope factor. Peak inward currents obtained from steady-state inactivation protocol were normalized by the maximum current amplitude and fit with a Boltzmann equation of the form  $I = I_{min} + (I_{max} - I_{min})/(1 + \exp[(V_m - V_{1/2})/k])$ , where  $V_m$  represents the inactivating prepulse membrane potential and  $V_{1/2}$  represents the midpoint of inactivation.

Recovery from inactivation was well described by a sum of two exponential components, and the development of inhibition in the presence of PF-05089771 was well described using a one-phase decay. A "plateau followed by a one phase decay" was employed for datasets where there was a steady-state baseline region before application of the compound. For experiments measuring the recovery from inactivation in the absence of PF-05089771, averaged data for each group was fit. For experiments measuring the development of inhibition in the presence of PF-05089771 and recovery from inhibition during washout of PF-05089771, individual cells were fit to provide the time constants, which were averaged and shown as mean  $\pm$  S.E.M.

## Results

**PF-05089771 Displays State-Dependent Inhibition of hNa<sub>v</sub>1.7 Currents.** PF-05089771 was previously shown to produce potent and selective inhibition of human Na<sub>v</sub>1.7 (IC<sub>50</sub> 11 nM) using a voltage protocol with a conditioning prepulse that evenly distributes the channel population across resting closed and inactivated states (Alexandrou et al., 2016). In the current study we examined in more detail the kinetics and state dependence of this interaction using the same protocol, which consists of an 8-second conditioning voltage step to an empirically determined membrane potential that results in 50% inactivation for each cell followed by 2-ms hyperpolarizing pulse to  $-120$  mV to partially relieve inactivation and then a 20-ms test pulse to 0 mV (Fig. 1A, right inset). This voltage protocol results in channels in both the fast and slow inactivated states that reach equilibrium during the progression of the pulse train (sweep interval of 15 seconds). At 100 nM PF-05089771, a concentration approximately ninefold higher than the previously reported IC<sub>50</sub>, hNa<sub>v</sub>1.7 currents are fully inhibited with a time constant of  $212 \pm 17$  seconds ( $n = 3$ , Figs. 1, A and B). The rate of inhibition at 1  $\mu$ M develops more rapidly, with a time constant of  $33 \pm 1$  seconds ( $n = 9$ ). Figures 1, A and B, also show that in the absence of an 8-second conditioning voltage step to promote inactivation, little or no inhibition during a 17-minute exposure to 100 nM PF-05089771 was observed, which is consistent with previously reported a lack of interaction with resting closed channels (Alexandrou et al., 2016).

To investigate the time course of PF-05089771 unblock that occurs when channels are returned to the resting closed state (i.e., at  $-120$  mV), we applied the half-inactivation protocol described above until maximal inhibition was established and then switched to a holding potential of  $-120$  mV, applying a



**Fig. 1.** Inhibition of human  $\text{Na}_v1.7$  by PF-05089771 develops and recovers slowly. (A) Representative traces showing the time course of inhibition under the three different recording conditions. Inhibition by PF-05089771 was assayed against resting channels (steady holding at  $-120$  mV) and a protocol in which channels were half-inactivated during 8-second prepulses delivered every 15 seconds from a steady holding voltage of  $-120$  mV ( $V_{0.5}$  of inactivation potential was determined empirically for each cell). The sweep interval for both protocols is 15 seconds. (B) Time course of inhibition for half-inactivated channels was dose dependent, reaching nearly complete block at both  $100$  nM ( $n = 3$ ) and  $1 \mu\text{M}$  ( $n = 9$ ); however, block was minimal for resting channels ( $n = 4$ ). (C) Recovery from block after washout of  $1 \mu\text{M}$  PF-05089771 using the tonic protocol (inset) is slow and incomplete ( $n = 5$ ).


20-ms test pulse every 10 seconds to assess magnitude of current recovery during washout of the compound. Recovery of current was slow ( $\tau = 526 \pm 36$  seconds,  $n = 5$ ) and incomplete, with current amplitude recovering slightly less than 60% of the predrug levels after 15 minutes (Fig. 1C). Failure to recover 100% of the current even after nearly 20 minutes may be partially due to rundown of the current over time.

**Using Brief Depolarizations, PF-05089771 Exhibits Very Slow Use-Dependent Block of  $\text{hNa}_v1.7$  Except at Very High Concentrations.** As the presence or absence of use-dependent block can provide a measure for the relative affinity for the open/fast-inactivated state, we assessed use-dependent inhibition by PF-05089771 using a 10-second high-frequency pulse train consisting of 100 steps (20 ms) to 0 mV at a rate of 10 Hz. As seen in Fig. 2A, we observed no difference between the DMSO control and 100 nM of PF-05089771, a concentration approximately ninefold higher than the published  $\text{IC}_{50}$  for partially inactivated channels (Alexandrou et al., 2016). Only at  $10 \mu\text{M}$  ( $\sim 900$ -fold over  $\text{IC}_{50}$ ) do we observe complete use-dependent block with a single 10 Hz train for 10 seconds. At concentrations less than  $10 \mu\text{M}$ , in excess of 100 pulses at 10 Hz was needed to achieve steady-state inhibition (data not shown). After complete block with  $10 \mu\text{M}$  PF-05089771, the currents were then allowed to recover in the absence of compound using the resting channel protocol (Fig. 2B). Currents recovered to at a similar rate ( $633 \pm 57$  seconds) but to a greater extent (about 75% of the predrug amplitude) within the same time frame as shown in Fig. 1C. The greater extent of current recovery may be partially explained by the  $\sim 6$  minutes shorter overall recording time (block and recovery) used for the experiment in Fig. 2, thus minimizing the contribution of current rundown.

The requirement of such high concentrations to observe use-dependent block could be attributable to a low affinity of PF-05089771 for the open/fast-inactivated state and/or the need to overcome the very slow apparent on-rate of the compound. In the following sections, we explored this question by testing the relative affinity for the fast- versus slow-inactivated states using voltage-clamp protocols that also account for the slow recovery from block of the compound.

**Investigating the Development of Inhibition for Fast- and Slow-Inactivated Channels.** Thus far the data presented in Figs. 1 and 2 show that the kinetics of and recovery from inhibition of  $\text{hNa}_v1.7$  by PF-05089771 are slow, with brief depolarizations (Fig. 2) resulting in no appreciable inhibition, even at a concentration ninefold higher than the  $\text{IC}_{50}$  for inactivated channels. However, these observations do not rule out the possibility of slow binding to fast-inactivated channels. In an effort to understand the mechanism of inhibition, we employed several voltage-clamp protocols that biased channels into predominantly fast- or slow-inactivated populations and subsequently measured the onset of inhibition by 100 nM PF-05089771.

We first determined the availability of fast- versus slow-inactivated states in the absence of PF-05089771 using inactivating step depolarizations to 0 mV for durations of 0.1, 1, 4, or 8 seconds followed by a hyperpolarizing step to  $-120$  mV for increasing durations to assess recovery from inactivation. As seen in Fig. 3A, the recovery from inactivation was well described by a sum of two exponential components with the shorter conditioning pulses increasing the relative availability of the fast-inactivated state (see Table 1). Not until the 4- and 8-second depolarizations do we observe a substantial fraction of the channels entering the slowly recovering state.




**Fig. 2.** By using brief depolarizations, PF-05089771 exhibits use dependence only at very high concentrations. (A) Use-dependent inhibition was examined by employing a standard high-frequency pulse train consisting of 100 steps (20 ms) to 0 mV at a rate of 10 Hz. The 10-Hz train was tested against the vehicle control (0.1% DMSO,  $n = 24$ ) and PF-05089771 at 100 nM ( $n = 5$ ), 1  $\mu$ M ( $n = 5$ ), 3  $\mu$ M ( $n = 5$ ), and 10  $\mu$ M ( $n = 9$ ). (B) After complete block with the 10-Hz train in the presence of 10  $\mu$ M PF-05089771, recovery was assessed using a tonic protocol (inset) while washing out the compound ( $n = 8$ ). The gray line is the curve fit from Fig. 1C, showcasing the similar rate but different extent of recovery from block.

To assess channel inhibition after either a short (1 second) or long step (8 seconds) to 0 mV, we used a fixed recovery interval of 1 and 3 seconds at  $-120$  mV, respectively. These recovery intervals allowed for a similar amount of current to be recovered ( $\sim 93\%$  available) in the absence of any compound ensuring that the current amplitude remained stable under control conditions for both protocols. Importantly, the 1- and

3-second recovery intervals were still sufficiently brief to not allow any appreciable recovery from PF-05089771-mediated inhibition. The compound was perfused until inhibition stabilized. As shown in Fig. 3C, plotting the fraction of available current over total time (time at 0 mV and  $-120$  mV), the block develops 4.5-fold faster for the 8-second pulse to 0 mV ( $\tau = 65 \pm 5$  seconds,  $n = 7$ ), compared with the 1-second pulse to 0 mV ( $\tau = 304 \pm 34$  seconds,  $n = 5$ ). If PF-05089771 interacts preferentially with the slow-inactivated state, this result could be explained by the higher fraction of channels in the slow-inactivated state (88%) after the 8-second step to 0 mV compared with the 1-second step (30%). However, this initial assessment does not take into account the fraction of the total time spent at rest when little or no block is expected to occur. Therefore, the data were replotted as the fraction of available current as a function of the total time spent at 0 mV (Fig. 3E). When reanalyzed in this manner the rate of block between both protocols was found to be indistinguishable ( $\tau = 31 \pm 1$  versus  $32 \pm 4$  seconds for the 8- and 1-second conditioning steps, respectively;  $P > 0.05$  by unpaired  $t$  test). Thus the apparent slower rate of block seen with the shorter conditioning pulse in Fig. 3C can be explained by the briefer availability (1 versus 8 seconds) of the depolarized state with which PF-05089771 can interact. These results suggest that the relative availability of slow versus fast-inactivated channels has less of an impact on the development of inhibition than the total time spent in the depolarized state.

In the second set of experiments to investigate whether PF-05089771 interacts more favorably with slow- or fast-inactivated channels, we employed two protocols in which the total protocol length and time spent at 0 mV before the test pulse are identical, but the fraction of channels populating the slow versus fast-inactivated states is different. This was achieved by applying a single conditioning voltage step to 0 mV for 4 seconds at a frequency of 0.1 Hz or a conditioning train of ten 400-ms steps to 0 mV at a frequency of 1 Hz. At both 0.1 and 1 Hz stimulation frequencies, the cell is depolarized to 0 mV for a total of 4 seconds before the test pulse (Fig. 4A, right). As shown in Fig. 4A, left, the recovery from inactivation was well described by a sum of two exponential components with the 0.1 Hz prepulse ( $\tau_{\text{fast}} = 9$  ms,  $A_{\text{fast}} = 28\%$ ;  $\tau_{\text{slow}} = 340$  ms,  $A_{\text{slow}} = 72\%$ ) providing greater availability of slow inactivated channels as compared with the 1-Hz prepulse ( $\tau_{\text{fast}} = 4$  ms,  $A_{\text{fast}} = 79\%$ ;  $\tau_{\text{slow}} = 162$  ms,  $A_{\text{slow}} = 21\%$ ). To assess block of the channel after the 0.1- or 1-Hz prepulse trains, a fixed recovery interval of 3 seconds at  $-120$  mV (Fig. 4B, right) was used for both conditioning protocols as this interval allowed for more than 90% of the current to be recovered in both protocols under control conditions. Again, because of the very slow rate of recovery from inhibition by PF-05089771, the 3-second recovery step between the conditioning pulse(s) and the test pulse is not sufficiently long to allow appreciable recovery from inhibition. As shown in Fig. 4B, left, the onset of inhibition is indistinguishable using the 1- or 0.1-Hz prepulse ( $\tau = 141 \pm 23$  and  $178 \pm 16$  second, respectively,  $n = 3$  each;  $P > 0.05$  by unpaired  $t$  test) despite the more than threefold difference in availability of slow inactivated channels. These results provide additional evidence that the relative availability of particular inactivated state conformations has little to no influence on the development of PF-05089771-mediated inhibition.



**Fig. 3.** Time course of block by PF-05089771 is independent of the availability of kinetically defined inactivated states. (A) Recovery from inactivation was assessed after a conditioning depolarizing pulse to 0 mV for 0.1, 1, 4, and 8 seconds in the absence of PF-05089771 ( $n = 4, 4, 4,$  and  $5,$  respectively). A variable recovery interval at  $-120$  mV followed each step to 0 mV. After the recovery interval, a 10-ms test pulse at 0 mV assessed the fraction of available current. As the conditioning pulse is increased from 0.1 to 8 seconds, a larger fraction of channels enter a slowly recovering state. (B) Individual traces from a single representative cell showing recovery from inactivation after either a 1-second (top traces) or an 8-second (bottom traces) pulse to 0 mV. The pulses represent the recovery intervals from 1 ms to 10 seconds in half-log intervals. (C) Onset of block by 100 nM of PF-05089771 was measured using a short (1 second,  $n = 5$ ) or long (8 seconds,  $n = 7$ ) inactivating conditioning pulse to 0 mV using the protocols shown in (D). The black dots overlaying the red data points represent the time-matched equivalent at 0 mV compared with blue data points (i.e., every 8 seconds). (D) A fixed recovery interval of 1 and 3 seconds at  $-120$  mV (which allows for  $>90\%$  of the current to recover) was used for the 1- and 8-second conditioning pulse, respectively. Sweep intervals of 10 and 20 seconds were used with the 1- and 8-second conditioning pulses, respectively. (E) Replotting as the amount of block as a function of the total time spent at 0 mV uncovers a similar time course of inhibition between the two protocols ( $\tau = 31 \pm 1$  versus  $32 \pm 4$  seconds for the 8- and 1-second conditioning steps, respectively;  $P > 0.05$  by unpaired  $t$  test).

In the final experiment to investigate whether PF-05089771 displays a preference for a particular inactivated state, we systematically varied the inactivating voltage while fixing the duration of the conditioning prepulse to 8 seconds. We set the conditioning voltage to potentials in which the channels are fully inactivated (0 and  $-60$  mV) or  $\sim 50\%$  inactivated ( $V_{0.5}$  of inactivation potential). The  $V_{0.5}$  of inactivation potential was determined empirically for each cell immediately before

testing by running a steady-state inactivation protocol (Fig. 5Ai). Across all cells tested, the average  $V_{0.5}$  of inactivation was  $-84 \pm 1$  mV, with  $\geq 99\%$  of the current inactivated at  $-60$  mV ( $n = 9$ , Fig. 5A). We first assessed the availability of fast- versus slow-inactivated states in the absence of PF-05089771 using an inactivating step depolarization to 0 mV,  $-60$  mV or the  $V_{0.5}$  potential for 8 seconds. As shown in Fig. 5B, left, the recovery from inactivation was well

TABLE 1

Best-fit values (for data shown in Fig. 3A) using a nonlinear regression with a sum of two exponentials

Tau values were calculated as  $1/K_{fast}$  and  $1/K_{slow}$ . The tau<sub>fast</sub> values were constrained to that of the fit for the 0.1-s conditioning pulse.

	0 mV for 0.1 s	0 mV for 1 s	0 mV for 4 s	0 mV for 8 s
A <sub>fast</sub> (%)	92	70	32	12
Tau <sub>fast</sub> (ms)	8	8	8	8
A <sub>slow</sub> (%)	8	30	68	88
Tau <sub>slow</sub> (ms)	108	326	566	580


described by a sum of two exponential components with the 8-second prepulse to 0 mV (from Table 1), providing greater availability of slow inactivated channels compared with the prepulse to -60 mV ( $\tau_{fast} = 11$  ms,  $A_{fast} = 83\%$ ;  $\tau_{slow} = 515$  ms,  $A_{slow} = 17\%$ ) or the V0.5 of inactivation ( $\tau_{fast} = 14$  ms,  $A_{fast} = 89\%$ ;  $\tau_{slow} = 308$  ms,  $A_{slow} = 11\%$ ). These results demonstrate that substantially less slow-inactivation is generated with the steps to -60 mV or the V0.5 potential consistent with previous observations for Na<sub>v</sub>1.3 (Jo and Bean, 2011).

To assess the rate of block by PF-05089771, a fixed recovery interval of 3 seconds at -120 mV followed the conditioning pulse and the total sweep duration (20 seconds) was equal for all three protocols. As shown in Fig. 5C, the time constant for the onset of block is similar between the conditioning pulse to 0 mV ( $\tau = 67 \pm 4$  seconds,  $n = 7$ ) and -60 mV ( $\tau = 90 \pm 7$  seconds,  $n = 5$ ;  $P > 0.05$  by ANOVA and Tukey's multiple comparison test), despite the strikingly different recovery


profiles seen in Fig. 5B. The rate of block was significantly slower using the conditioning pulse to the V0.5 of inactivation ( $\tau = 331 \pm 56$  seconds,  $n = 6$ ) than that at either 0 or -60 mV ( $P < 0.01$  by ANOVA and Tukey's multiple comparison test). These results provide further evidence that PF-05089771 interacts rather indiscriminately with fast- or slow-inactivated channels and gives further indication that there is little or no interaction with resting channels. Additionally, inactivation at 0 mV proceeds primarily through the open state, whereas inactivation at -60 mV proceeds primarily through the closed state, as demonstrated by a lack of inward current generated with a step depolarization to -60 mV (Fig. 5A). Therefore, the results in Fig. 5C suggest that the rate and magnitude of inhibition is largely independent of the route of entry into the inactivated state, as well as the kinetically defined inactivated state in which the channel resides. The single commonality between the various protocols to observe channel inhibition is the requirement for a sufficiently strong depolarization to move the channels out of the resting conformation.

## Discussion

We previously reported two human Na<sub>v</sub>1.7 selective aryl sulfonamide small molecule inhibitors, PF-04856264 (McCormack et al., 2013) and PF-05089771 (Alexandrou et al., 2016) which exhibit strong state dependence, preferentially inhibiting inactivated channels with  $\geq 1000$ -fold higher



**Fig. 4.** Onset of inhibition develops over similar time course using different conditioning trains with equal time at 0 mV. (A) Recovery from inactivation was assessed after a conditioning prepulse train to 0 mV at a frequency of 1 Hz ( $n = 5$ ) and 0.1 Hz ( $n = 4$ ) in the absence of PF-05089771. (B) Onset of block by 100 nM of PF-05089771 ( $n = 3$ , each) was measured using both protocols but with a fixed recovery interval of 3 seconds at -120 mV and a sweep interval of 20 seconds. The onset of block is similar with both protocols ( $\tau = 141 \pm 23$  seconds for the  $10 \times 400$  ms, 1 Hz and  $178 \pm 16$  seconds for the  $1 \times 4$  seconds, 0.1 Hz,  $n = 3$  each;  $P > 0.05$  by unpaired  $t$  test), despite the  $\sim 3.4$ -fold difference in relative availability of slow inactivated channels with the 0.1-Hz conditioning stimulus.



**Fig. 5.** Onset of inhibition develops over similar time course from closed- and open-state inactivation. (A) Activation and steady-state inactivation curve show minimal overlap. The voltage dependence of steady-state inactivation (circles) was determined using the protocol shown in (i) with a  $V_{0.5}$  of  $-83.5 \pm 0.5$  mV ( $n = 9$ ). The voltage-dependence of activation (squares) was determined using the protocol shown in (ii) with a  $V_{0.5}$  of  $-29.3 \pm 0.3$  mV ( $n = 11$ ). (B) Recovery from inactivation was assessed after a conditioning pulse at 0 mV (channels inactivated through open state,  $n = 5$ ),  $-60$  mV (channels predominantly inactivated through closed state,  $n = 3$ ) or the  $V_{0.5}$  of inactivation (as determined empirically for each cell,  $n = 3$ ). For the conditioning pulses to  $-60$  and 0 mV, 100% of the channels are inactivated. The rate of recovery for the  $-60$  mV and  $V_{0.5}$  conditioning steps have a considerably larger fast component compared with the conditioning step to 0 mV. Likewise, the recovery from the  $V_{0.5}$  conditioning pulse is predominantly fast. (C) Onset of block by 100 nM of PF-05089771 was measured using each protocol with a fixed 3-second recovery interval at  $-120$  mV and a 20-second sweep interval. The rate of block is similar for the 0 mV ( $n = 7$ ) and  $-60$  mV ( $n = 5$ ) protocols ( $P > 0.05$  by ANOVA), despite having dramatically different recovery profiles; however, the onset of block is dramatically slower using the  $V_{0.5}$  potential ( $P < 0.01$  by ANOVA and Tukey's multiple comparison test,  $n = 6$ ).

$IC_{50}$  compared with resting channels. In addition to the pronounced state dependence and  $Na_v$  selectivity an unexplored attribute of these molecules is the apparent slow association and disassociation with  $Na_v1.7$ . In the present study, we sought to further investigate whether the pronounced degree of state-dependent inhibition by PF-05089771 could be attributed to a preference for a particular inactivated state of the channel.

Commonly used two-pulse protocols that examine the voltage or time dependence of fast- versus slow-inactivation cannot reliably differentiate compounds that bind preferentially to slow-inactivated state(s) versus those that associate slowly to inactivated channels, particularly if a compound is present while the channels transition through multiple inactivated states as during a long depolarizing pulse (Lenkey et al., 2006; Karoly et al., 2010). Typically these

protocols measure the steady-state shifts in availability of slow- and fast-inactivated states in the presence of a state-dependent blocker. One prerequisite of such protocols is the need to "reset" the system before each successive change in conditioning pulse voltage or duration to avoid the accumulation of block in successive sweeps. The slow onset of inhibition with PF-05089771 coupled with the slow and incomplete washout makes use of such protocols unfeasible. We therefore modified the voltage protocols to take advantage of the unique properties of PF-05089771 and allow more quantitative characterization of PF-05089771-mediated inhibition. Using these protocols it was possible to increase availability of the fast-inactivated state without generating appreciable slow inactivation. The slow time course of unblock of PF-05089771 minimized recovery from inhibition between successive sweeps and provided long temporal access (within

each sweep) to primarily fast-inactivated channels, thereby allowing us to discern slow binding to a population of channels that are predominantly fast inactivated.

The results presented here suggest that for the development of PF-05089771-mediated inhibition, the preference for a particular inactivated state (fast and/or slow) appears less critical than the relative time that the channel is depolarized. When accounting for the time that the channel is depolarized, the rate of inhibition by PF-05089771 occurs with a similar time course for channels that reside in either predominantly fast- or slow-inactivated states. Thus the presentation of a particular inactivated state does not appear to be a prerequisite for association of PF-05089771 with the channel. Likewise, these observations do not posit that PF-05089771 interacts with only fast-inactivated channels, as gating changes that occur with long depolarizations or at more depolarized potentials (both of which are associated with the development of slow-inactivation) do not impede inhibition. As such, we propose that the only requirement for PF-05089771-mediated inhibition is a sufficiently strong depolarization to move the channel out of the resting/closed configuration (for which there is very low affinity). Any subsequent conformational changes associated with fast- or slow-inactivation appear to have little observable effect on the interaction of PF-05089771 with the channel. A caveat to this interpretation is that we are unable to employ a protocol that allows a particular inactivated state to be assessed in isolation, because the assumption that channels populate a single fast- or single slow-inactivated state is an oversimplification, and it is likely that the channel moves through a series of intermediate conformational changes during the gating process. Furthermore, it has been proposed that that fast- and slow-inactivation are not mutually exclusive, representing structurally independent, nonsequential processes with both states potentially existing in the same channel simultaneously (Bezanilla et al., 1982; VEDANTHAM and Cannon, 1998).

Utilizing mutagenesis studies, we previously demonstrated that Na<sub>v</sub>1.7-selective aryl sulfonamide inhibitors interact with the extracellular surface of the VSD region of DIV (McCormack et al., 2013; Alexandrou et al., 2016), although the precise mechanism for inhibition was not fully understood. Over the years, the use of polypeptide toxins has increased our understanding of the functional significance of the VSD of DIV in channel fast-inactivation (Catterall et al., 2007; Hanck and Sheets, 2007; Groome, 2014). Site-3 toxins, which include  $\alpha$ -scorpion toxins and sea anemone toxins, bind to overlapping extracellular residues on the DIV S3–S4 loop, binding with highest affinity at negative membrane potentials where the channel is closed (Catterall et al., 2007). These site-3 toxins impede fast inactivation by trapping the DIV S4 segment and preventing outward translocation upon depolarization (Rogers et al., 1996; Xiao et al., 2014). At least one of the residues (D1586) of the proposed PF-05089771 binding site, located near the extracellular end of the S3 segment, overlaps with the binding site for site-3 toxins. Thus it remains a possibility that PF-05089771 employs a similar voltage-sensor trapping mechanism to stabilize a nonconducting conformation of the channel. As it has been suggested, that movement of the DIV voltage-sensor after depolarization is necessary and sufficient for inactivation via the open (i.e., strong depolarization to 0 mV) or closed states (i.e., mild depolarization to –60 mV) (Capes et al., 2013), we hypothesize that translocation

of the DIV voltage-sensor upon depolarization exposes the high-affinity small molecule binding site and binding of PF-05089771 stabilizes the channel into a nonconducting conformation from which recovery to a resting conformation is slow. A recent study by Ahuja et al. (2015) provides strong support for this hypothesis. Using a chimeric construct of human Na<sub>v</sub>1.7 DIV VSD and the bacterial Na<sub>v</sub>Ab, they were able to demonstrate through both electrophysiological and crystallographic methods that Na<sub>v</sub>1.7 aryl sulfonamide inhibitors employ a voltage-sensor trapping mechanism to lock the activated DIV voltage-sensor into the activated conformation and thereby stabilizing a nonconductive conformation of the channel.

Antiepileptics and anticonvulsants provide therapeutic utility via use-dependent inhibition of fast firing neurons typically associated with epileptic seizures and pain (Cummins and Rush, 2007). Alternatively, as sodium channels undergo slow inactivation after sustained depolarizations, neurons that are abnormally depolarized for long periods of time (after injury or insult) may be susceptible to enhanced block via slow-inactivation preferring drugs (Errington et al., 2008; Hildebrand et al., 2011; Jo and Bean, 2011). As PF-05089771 and its congeners interact with the VSD to inhibit depolarized human Na<sub>v</sub>1.7 channels, irrespective of the channel state, this mechanism of action may provide a therapeutic benefit across multiple types of pain modalities characterized by any type of Na<sub>v</sub>1.7 channel activity. We believe that the development of small molecules that act as Na<sub>v</sub> channel gating modifiers via an interaction with the VSD, as opposed to more classic nonselective inhibitors that bind inside the channel pore, represents an avenue worthy of further exploration in an effort to obtain improved efficacy and isoform selectivity.

#### Acknowledgments

The authors thank Sally Stoehr for help with initial experiments, Eva Prazak and Doug McIlvaine for providing cells, and Neil Castle and Doug Krafte for suggestions during the drafting of the manuscript.

#### Authorship contributions

*Participated in research design:* Theile, Fuller, and Chapman.  
*Conducted experiments:* Theile and Fuller.  
*Performed data analysis:* Theile, Fuller, and Chapman.  
*Wrote or contributed to the writing of the manuscript:* Theile and Chapman.

#### References

- Ahuja S, Mukund S, Deng L, Khakh K, Chang E, Ho H, Shriver S, Young C, Lin S, Johnson JP, Jr, et al. (2015) Structural basis of Nav1.7 inhibition by an isoform-selective small-molecule antagonist. *Science* **350**:aac5464.
- Alexandrou AJ, Brown AR, Chapman ML, Estacion M, Turner J, Mis MA, Wilbrey A, Payne EC, Gutteridge A, Cox PJ, et al. (2016) Subtype-selective small molecule inhibitors reveal a fundamental role for Nav1.7 in nociceptor electrogenesis, axonal conduction and presynaptic release. *PLoS One* **11**:e0152405.
- Bezanilla F, Taylor RE, and Fernández JM (1982) Distribution and kinetics of membrane dielectric polarization. I. Long-term inactivation of gating currents. *J Gen Physiol* **79**:21–40.
- Capes DL, Goldschen-Ohm MP, Arcisio-Miranda M, Bezanilla F, and Chanda B (2013) Domain IV voltage-sensor movement is both sufficient and rate limiting for fast inactivation in sodium channels. *J Gen Physiol* **142**:101–112.
- Catterall WA (2000) From ionic currents to molecular mechanisms: the structure and function of voltage-gated sodium channels. *Neuron* **26**:13–25.
- Catterall WA (2010) Ion channel voltage sensors: structure, function, and pathophysiology. *Neuron* **67**:915–928.
- Catterall WA, Cestèle S, Yarov-Yarovsky V, Yu FH, Konoki K, and Scheuer T (2007) Voltage-gated ion channels and gating modifier toxins. *Toxicol* **49**:124–141.
- Cox JJ, Reimann F, Nicholas AK, Thornton G, Roberts E, Springell K, Karbani G, Jafri H, Mannan J, Raashid Y, et al. (2006) An SCN9A channelopathy causes congenital inability to experience pain. *Nature* **444**:894–898.



- Cummins TR and Rush AM (2007) Voltage-gated sodium channel blockers for the treatment of neuropathic pain. *Expert Rev Neurother* **7**:1597–1612.
- Dib-Hajj SD, Black JA, and Waxman SG (2009) Voltage-gated sodium channels: therapeutic targets for pain. *Pain Med* **10**:1260–1269.
- Dib-Hajj SD, Yang Y, and Waxman SG (2008) Genetics and molecular pathophysiology of Na(v)1.7-related pain syndromes. *Adv Genet* **63**:85–110.
- Errington AC, Stöhr T, Heers C, and Lees G (2008) The investigational anticonvulsant lacosamide selectively enhances slow inactivation of voltage-gated sodium channels. *Mol Pharmacol* **73**:157–169.
- Fozzard HA, Lee PJ, and Lipkind GM (2005) Mechanism of local anesthetic drug action on voltage-gated sodium channels. *Curr Pharm Des* **11**:2671–2686.
- Groome JR (2014) The voltage sensor module in sodium channels. *Handbook Exp Pharmacol* **221**:7–31.
- Hanck DA and Sheets MF (2007) Site-3 toxins and cardiac sodium channels. *Toxicol* **49**:181–193.
- Hildebrand ME, Smith PL, Bladen C, Eduljee C, Xie JY, Chen L, Fee-Maki M, Doering CJ, Mezeyova J, Zhu Y, et al. (2011) A novel slow-inactivation-specific ion channel modulator attenuates neuropathic pain. *Pain* **152**:833–843.
- Hodgkin AL and Huxley AF (1952) A quantitative description of membrane current and its application to conduction and excitation in nerve. *J Physiol* **117**:500–544.
- Jo S and Bean BP (2011) Inhibition of neuronal voltage-gated sodium channels by brilliant blue G. *Mol Pharmacol* **80**:247–257.
- Karoly R, Lenkey N, Juhász AO, Vizi ES, and Mike A (2010) Fast- or slow-inactivated state preference of Na<sup>+</sup> channel inhibitors: a simulation and experimental study. *PLOS Comput Biol* **6**:e1000818.
- Lenkey N, Karoly R, Kiss JP, Szasz BK, Vizi ES, and Mike A (2006) The mechanism of activity-dependent sodium channel inhibition by the antidepressants fluoxetine and desipramine. *Mol Pharmacol* **70**:2052–2063.
- McCormack K, Santos S, Chapman ML, Krafte DS, Marron BE, West CW, Krambis MJ, Antonio BM, Zellmer SG, Printzenhoff D, et al. (2013) Voltage sensor interaction site for selective small molecule inhibitors of voltage-gated sodium channels. *Proc Natl Acad Sci USA* **110**:E2724–E2732.
- Noda M, Shimizu S, Tanabe T, Takai T, Kayano T, Ikeda T, Takahashi H, Nakayama H, Kanaoka Y, Minamino N, et al. (1984) Primary structure of Electrophorus electricus sodium channel deduced from cDNA sequence. *Nature* **312**:121–127.
- Payne CE, Brown AR, Theile JW, Loucif AJ, Alexandrou AJ, Fuller MD, Mahoney JH, Antonio BM, Gerlach AC, Printzenhoff DM, et al. (2015) A novel selective and orally bioavailable Nav 1.8 channel blocker, PF-01247324, attenuates nociception and sensory neuron excitability. *Br J Pharmacol* **172**:2654–2670.
- Ragsdale DS, McPhee JC, Scheuer T, and Catterall WA (1994) Molecular determinants of state-dependent block of Na<sup>+</sup> channels by local anesthetics. *Science* **265**:1724–1728.
- Ragsdale DS, McPhee JC, Scheuer T, and Catterall WA (1996) Common molecular determinants of local anesthetic, antiarrhythmic, and anticonvulsant block of voltage-gated Na<sup>+</sup> channels. *Proc Natl Acad Sci USA* **93**:9270–9275.
- Rogers JC, Qu Y, Tanada TN, Scheuer T, and Catterall WA (1996) Molecular determinants of high affinity binding of alpha-scorpion toxin and sea anemone toxin in the S3-S4 extracellular loop in domain IV of the Na<sup>+</sup> channel alpha subunit. *J Biol Chem* **271**:15950–15962.
- Stühmer W, Conti F, Suzuki H, Wang XD, Noda M, Yahagi N, Kubo H, and Numa S (1989) Structural parts involved in activation and inactivation of the sodium channel. *Nature* **339**:597–603.
- Theile JW and Cummins TR (2011) Recent developments regarding voltage-gated sodium channel blockers for the treatment of inherited and acquired neuropathic pain syndromes. *Front Pharmacol* **2**:1–14.
- Vedantham V and Cannon SC (1998) Slow inactivation does not affect movement of the fast inactivation gate in voltage-gated Na<sup>+</sup> channels. *J Gen Physiol* **111**:83–93.
- Wang J, Yarov-Yarovoy V, Kahn R, Gordon D, Gurevitz M, Scheuer T, and Catterall WA (2011) Mapping the receptor site for alpha-scorpion toxins on a Na<sup>+</sup> channel voltage sensor. *Proc Natl Acad Sci USA* **108**:15426–15431.
- Xiao Y, Blumenthal K, and Cummins TR (2014) Gating-pore currents demonstrate selective and specific modulation of individual sodium channel voltage-sensors by biological toxins. *Mol Pharmacol* **86**:159–167.
- Yarov-Yarovoy V, Brown J, Sharp EM, Clare JJ, Scheuer T, and Catterall WA (2001) Molecular determinants of voltage-dependent gating and binding of pore-blocking drugs in transmembrane segment IIIIS6 of the Na<sup>(+)</sup> channel alpha subunit. *J Biol Chem* **276**:20–27.
- Yarov-Yarovoy V, McPhee JC, Idsvoog D, Pate C, Scheuer T, and Catterall WA (2002) Role of amino acid residues in transmembrane segments IS6 and IIS6 of the Na<sup>+</sup> channel alpha subunit in voltage-dependent gating and drug block. *J Biol Chem* **277**:35393–35401.

---

**Address correspondence to:** Jonathan Theile, Icagen, Inc., 4222 Emperor Boulevard, Suite 350, Durham, NC 27703. E-mail: jtheile@icagen.com

---